

Glueball- $Q\bar{Q}$ mixing and Okuba-Zweig-Lizuka rule violation in the hadronic decays of heavy quarkonia

Qiang Zhao^{1,2*}, Bing-Song Zou^{1†}, and Zhong-Biao Ma³

1) *Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, 100049, P.R. China*

2) *Department of Physics, University of Surrey, Guildford, GU2 7XH, United Kingdom and*

3) *CCAST, Beijing 100080, P.R. China*

(Dated: February 2, 2008)

We investigate the correlations between the scalar meson configurations and Okuba-Zweig-Lizuka (OZI) rule violations in the hadronic decays of heavy quarkonia, e.g. J/ψ and Υ , into isoscalar vector meson (ϕ and ω) and scalar mesons ($f_0(1710)$, $f_0(1500)$, and $f_0(1370)$). It shows that the dramatic change of the values of the branching ratio fraction of $\phi f_0/\omega f_0$ from low (e.g. in J/ψ decays) to high energies (e.g. in Υ decays) will not only test the glueball- $Q\bar{Q}$ mixings, but also provide important information about the mysterious OZI-rule violations within the scalars.

The recent experimental data from BES [1, 2, 3, 4] seem to have enhanced the possible existence of significant glueball contents within the low-lying scalars [5, 6, 7, 8, 9, 10, 11, 12, 13]. In particular, large glueball components are preferred in the $f_0(1500)$, while large $|s\bar{s}\rangle$ and $|n\bar{n}\rangle \equiv |u\bar{u} + d\bar{d}\rangle/\sqrt{2}$ are dominant in the $f_0(1710)$ and $f_0(1370)$, respectively. Although the glueball components are relatively small in these two states, their phases relative to the dominant $Q\bar{Q}$ components turn out to play crucial roles in the understanding of the scalar spectroscopy [5, 6, 14, 15]. Implications arising from the presence of the glueball components can then be detected in their production processes. Mechanisms such as J/ψ hadronic and radiative decays are ideal for probing information about the glueball- $Q\bar{Q}$ mixings.

In Ref. [15], we reported a coherent study of the production of f_0^i ($i = 1, 2, 3$ corresponding to $f_0(1710)$, $f_0(1500)$, and $f_0(1370)$) in $J/\psi \rightarrow V f_0 \rightarrow VPP$, where $V = \phi, \omega$. There are advantages of studying isoscalar vector meson ϕ and ω in association with the f_0 states: i) Since ω and ϕ are approximately ideally mixed in the flavor space, the process $J/\psi \rightarrow \phi f_0(s\bar{s})$ and $J/\psi \rightarrow \omega f_0(n\bar{n})$ may probe the $s\bar{s}$ and $n\bar{n}$ components separately to leading order via the singly OZI [16] disconnected diagrams (Fig. 1(b)); ii) If there exist glueball components in the f_0 states, those two processes also probe the glueball components at the same order via $J/\psi \rightarrow VG$ (Fig. 1(a)); iii) As discussed in Ref. [15], the ratio of these two decay channels also highlights the importance of the doubly OZI disconnected diagram in the interpretation of the BES data for the production of $f_0(1370)$ and $f_0(1500)$ in the J/ψ decays (see Fig. 1(c)), which is contrary to expectations based on naive OZI rule [17]. Therefore, the importance of its violations can also be reflected by the failure of the prescriptions, which only include the leading contributions from the singly disconnected processes and the direct glueball production.

It was pointed out by Lipkin before that the OZI-rule violations can always proceed by a two-step process involving intermediate virtual mesons [18]. For example, as studied by Geiger and Isgur [19], systematic cancellations among the hadronic loops occurred for the $u\bar{u} \leftrightarrow s\bar{s}$ mixing in all the low-lying nonets except 0^{++} . A general argument was given by Lipkin and Zou [20]. Similar to the large rates of $p\bar{p} \rightarrow K^*\bar{K} + c.c. \rightarrow \phi\pi$ [21], the doubly OZI disconnected processes in the J/ψ decays, can occur via $J/\psi \rightarrow K^*\bar{K} + c.c. \rightarrow V f_0^i$, or $J/\psi \rightarrow \rho\pi + c.c. \rightarrow V f_0^i$. As we know that $J/\psi \rightarrow K^*\bar{K} + c.c.$ and $\rho\pi + c.c.$ are two of the largest decay channels of J/ψ [22], the virtual meson mediated transitions as the dominant contributions to the doubly OZI disconnected processes may not be small. Such an approach was applied to the investigation of the “ $\rho\pi$ puzzle” in J/ψ and ψ' by Li, Bugg and Zou [23]. In comparison with the quark-gluon interaction picture the intermediate meson exchange model could be useful for us to study the correlation between the strong glueball- $Q\bar{Q}$ mixings and presence of possible OZI violations. In particular, the energy evolution of the doubly disconnected processes can be investigated and may shed light on some of those recently raised puzzles in the scalar meson productions [4].

As follows, we first summarize the glueball- $Q\bar{Q}$ mixing scheme and the factorizations for the f_0^i productions in the quarkonium decays. By studying the exclusive reactions, $J/\psi \rightarrow V f_0^i \rightarrow VPP$, and $\Upsilon \rightarrow V f_0^i \rightarrow VPP$, where the “close-to-ideally-mixed” ω and ϕ play a role as an OZI filter, the different energy-dependence between the singly and doubly disconnected processes can be highlighted. We then study the energy evolution of the doubly OZI disconnected processes in an intermediate meson exchange model. The conclusion will be drawn in the end.

Applying the glueball- $Q\bar{Q}$ mixing scheme proposed by Amsler and Close [6], and developed later by Close *et*

* e-mail: qiang.zhao@surrey.ac.uk

† e-mail: zoubs@ihep.ac.cn

al [14, 15], the three f_0^i states, which are the eigenstates of the glueball- $Q\bar{Q}$ mixing potential, can be described as

$$\begin{pmatrix} |f_0(1710)\rangle \\ |f_0(1500)\rangle \\ |f_0(1370)\rangle \end{pmatrix} = \begin{pmatrix} x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \\ x_3 & y_3 & z_3 \end{pmatrix} \begin{pmatrix} |G\rangle \\ |s\bar{s}\rangle \\ |n\bar{n}\rangle \end{pmatrix}, \quad (1)$$

where $|G\rangle$, $|s\bar{s}\rangle$, and $|n\bar{n}\rangle$ are the pure glueball and $Q\bar{Q}$ nonet, respectively, and x_i , y_i and z_i are the mixing matrix elements determined by the flavor-independent glueball- $Q\bar{Q}$ transitions [6, 14]. As studied in Ref. [14, 15], the mixing matrix determined by the data for $f_0^i \rightarrow PP$ possesses stable features and can be tested in the f_0^i production processes. In particular, it was shown in Ref. [15] that the description of $J/\psi \rightarrow Vf_0^i$ needs essentially a strong glueball component in the $f_0(1500)$. Furthermore, the preference of the presence of strong glueball component in the f_0^i also implies the strong OZI violations could become significant at the low energy region.

In Ref. [15], an OZI-selecting factorization scheme is proposed for $J/\psi \rightarrow Vf_0^i$, and can be generalized to other vector quarkonium decays, e.g. $\Upsilon \rightarrow Vf_0^i$. The transition occurs via

$$\begin{aligned} M_{\phi G} &\equiv \langle G|V_\phi|Q_V\rangle \\ M_{\phi(s\bar{s})} &\equiv \langle s\bar{s}|V_\phi|Q_V\rangle \\ M_{\phi(n\bar{n})} &\equiv \langle n\bar{n}|V_\phi|Q_V\rangle, \end{aligned} \quad (2)$$

where Q_V denotes the quarkonium wavefunction, and $M_{\phi G}$, $M_{\phi(s\bar{s})}$, and $M_{\phi(n\bar{n})}$ are transition amplitudes for the production of the pure glueball and flavor nonet through the vector meson potential. Since the ϕ is almost $s\bar{s}$, the potential filters out the higher order OZI process $M_{\phi(n\bar{n})}$, which is doubly OZI disconnected at leading order.

Due to lack of information about the potentials, we assume, based on the order of α_s , that the singly OZI disconnected processes have the same strengths. As an example for $J/\psi \rightarrow \phi f_0^i$, we assume

$$\langle G|V_\phi|J/\psi\rangle \simeq \langle s\bar{s}|V_\phi|J/\psi\rangle. \quad (3)$$

For the doubly disconnected transition $M_{\phi(n\bar{n})}$, it is natural to introduce r for its relative strength to the singly disconnected ones:

$$\langle n\bar{n}|V_\phi|J/\psi\rangle \simeq \sqrt{2}r\langle s\bar{s}|V_\phi|J/\psi\rangle, \quad (4)$$

where the factor $\sqrt{2}$ is from the normalization of $n\bar{n}$. Naively, $r \ll 1$ would imply that the doubly disconnected process has a perturbative feature, while $r \simeq 1$ reflects the strong OZI-rule violation. The same factorization can also be applied to $J/\psi \rightarrow \omega f_0^i$ [15].

Following the assumption made for those three independent transitions, and considering the partial decay widths for $J/\psi \rightarrow \phi f_0^i \rightarrow \phi PP$ and $J/\psi \rightarrow \omega f_0^i \rightarrow \omega PP$, we have a set of equations in terms of only one independent transition amplitude of the pure glueball production:

$$\Gamma_{J/\psi \rightarrow \phi f_0^i \rightarrow \phi PP} = \frac{|\mathbf{p}_{\phi i}|}{|\mathbf{p}_{\phi G}|} b r_{f_0^i \rightarrow PP} [x_i + y_i + \sqrt{2}r z_i]^2 \Gamma_{J/\psi \rightarrow \phi G}, \quad (5)$$

$$\Gamma_{J/\psi \rightarrow \omega f_0^i \rightarrow \omega PP} = \frac{|\mathbf{p}_{\omega i}|}{|\mathbf{p}_{\omega G}|} b r_{f_0^i \rightarrow PP} [x_i + r y_i + \sqrt{2}z_i]^2 \Gamma_{J/\psi \rightarrow \omega G}, \quad (6)$$

where $2\Gamma_{J/\psi \rightarrow \phi G}/|\mathbf{p}_{\phi G}| = \Gamma_{J/\psi \rightarrow \omega G}/|\mathbf{p}_{\omega G}|$ due to the flavor-blind assumption. For f_0^i decays into the same pseudoscalar meson pairs, such as $f_0^i \rightarrow K\bar{K}$, the ratio between these two partial decay widths becomes

$$R_i^{OZI} = \frac{\Gamma_{J/\psi \rightarrow \phi f_0^i \rightarrow \phi K\bar{K}}}{\Gamma_{J/\psi \rightarrow \omega f_0^i \rightarrow \omega K\bar{K}}} = \frac{|\mathbf{p}_{\phi i}|}{|\mathbf{p}_{\omega i}|} \frac{[x_i + y_i + \sqrt{2}r z_i]^2}{2[x_i + r y_i + \sqrt{2}z_i]^2}, \quad (7)$$

which will be the focus of our discussions in this work. First, we point out that, apart from the momenta, $|\mathbf{p}_{\phi i}|$ and $|\mathbf{p}_{\omega i}|$, the energy scale does not explicitly appear in the above expression. The ratio R_i^{OZI} will then be determined correlatively by the scalar mixing matrix elements and the doubly OZI violations (r). Also, we note that the mixing matrix elements contain the structure information about the scalars and are independent of the energy scales. In contrast, the doubly OZI violation parameter r could depend on the energy scale. Therefore, it is natural to expect that the energy-dependence of r would lead to the change of R_i^{OZI} at different decay energies, or between, e.g., J/ψ and Υ decays. Also, due to the correlation between r and the mixing matrix elements, certain pattern for the energy-dependence of R_i^{OZI} can be predicted.

As an example, to the limit of $r \rightarrow 0$, where the OZI rule applies, Eq. (7) for the $f_0(1710)$ production will reduce to

$$R_1^{OZI} \simeq \frac{|\mathbf{p}_{\phi 1}|}{2|\mathbf{p}_{\omega 1}|} \frac{[x_1 + y_1]^2}{[x_1 + \sqrt{2}z_1]^2} \simeq 3. \quad (8)$$

In the case of J/ψ decays [15], this value is contradicted by the experimental data $R_1^{exp} \simeq 0.15$ from BES [1, 2], and 0.75 from PDG [22]. Substituting the experimental data into the above equation, one determines $r \simeq 2.2$ and 0.5, respectively. This suggests that there are important contributions from the OZI rule violations in the J/ψ decays.

Although it may not be very surprising that there are significant OZI violation contributions in the scalar productions, it nontrivially implies that its presence and absence would lead to inversely changed values for R_1^{OZI} as shown by Eqs. (7) and (8). Note that though Eq. (7) is derived in the J/ψ decays, it does not strongly depend on the masses of the J/ψ meson except for r , of which the energy-dependence is not known [27]. In another word, by measuring the energy dependence of

$$R_i^{OZI} \equiv \frac{\Gamma_{3g(1S) \rightarrow \phi f_0^i}}{\Gamma_{3g(1S) \rightarrow \omega f_0^i}} \simeq \frac{[x_i + y_i + \sqrt{2}rz_i]^2}{2[x_i + ry_i + \sqrt{2}z_i]^2}, \quad (9)$$

one would have access to the energy evolution of the OZI violation contributions contained within r . As a result, the correlated information about the scalar structures can also be examined.

Equation (9) allows us to investigate the evolution of R_i^{OZI} in terms of a range of r , e.g. $r = 0 \sim 3$. In such a general context, $r \rightarrow 0$ will correspond to the regime where the OZI rule is well respected, while $r \sim 1$, to the region of strong OZI violations. We plot the ratio changes in Fig. 2. For the J/ψ decays, $r \simeq 2.2$ suggests the significant breaking of the OZI rule as found in Ref. [15]. This is the region that the small fraction of $R_1^{OZI} = 0.15$ for the $f_0(1710)$ is observed. Furthermore, the ratios R_2^{OZI} and R_3^{OZI} for the $f_0(1500)$ and $f_0(1370)$ are much larger than unit. Hence, the branching ratio pattern observed by BES emerges.

In Fig. 2, in the limit of $r \rightarrow 0$, R_i^{OZI} exhibits dramatical changes. In particular, it shows that the production branching ratio for $\phi f_0(1710)$ will become much larger than that for $\omega f_0(1710)$. Therefore, the ratio $R_1^{OZI} = 0.15$ observed in the J/ψ decays will be inversely changed to $R_1^{OZI} > 1$. In fact, such an inverse change could happen even at moderate values for r . As shown by Fig. 2, the ratio $R_1^{OZI} = 1$ corresponds to $r = 0.5$. In brief, Fig. 2 provides a guide for the change of the branching ratio fractions at different energies via the energy evolution of r . However, such an evolution trend will only become useful when the energy dependence of r is quantified. This requires an explicit relation between the doubly and singly OZI disconnected processes to be established. It will then enable us to map r in Fig. 2 to a certain energy region, where the ratio R_i^{OZI} can be predicted.

Theoretically, the energy evolution of r can be investigated by calculating the relative strengths between the doubly disconnected process and the singly disconnected one as defined by Eq. (4). However, due to the break-down of the perturbative character of the doubly disconnected processes, Fig. 1(c) cannot be regarded as Feynman diagram, but rather a topological one. In this sense, a pQCD calculation of r becomes unreliable. We hence introduce a dynamical assumption based on successful phenomenological treatments available in the literature. We assume that the OZI doubly disconnected processes are dual with the intermediate meson exchange processes [18, 19, 20].

In the picture of Geiger and Isgur's intermediate virtual meson exchanges, the doubly disconnected diagram can be illustrated by two topological diagrams, Fig. 3(a) and (b). Qualitatively, such processes will be suppressed relative to the singly disconnected ones at higher energies. In Fig. 3(a), the two colored quark-antiquark pairs must exchange a colored gluon to keep overall color-singlet. In the case that the created quark pairs scatter out to form the final state V and f_0^i , there must need large momentum transfers between quark line q_1 and q_2 . Nevertheless, the momentum difference between q_1 and q_2 will increase with the increasing masses of the initial quarkonia since they must be recoiled back to the directions of their original quark (antiquark) partners to form the mesons in the final state. As a consequence, the exchanged gluon will become very hard and the transition amplitude will be suppressed at higher masses. One can also understand this as the suppression from the wavefunction distribution of the virtual meson formed by q_1 and q_2 . Higher energies will correspond to the wavefunction tail at shorter distances, which is far away from the large wavefunction density determined by the typical size of meson.

In Fig. 3(b), the color-singlet quark-antiquark pair of the scalar meson is created and recoiled to the backward direction. In the virtual meson exchange model, as shown by Fig. 3(b), the quark-antiquark pairs must correlate with each other to form the virtual mesons. Therefore, at least at one vertex, e.g. V_q , which is the $^3P_0$ production vertex, the quark line must be recoiled back approximately to the momentum direction of the final antiquark. Because of such a large momentum transfer at vertex V_q , the $^3P_0$ mechanism is suppressed, which means that the transition via Fig. 3(b) will decrease with the increasing energies, i.e. in the decay of heavier quarkonia.

In principle, by summing over all the possible meson exchange loops, the meson exchange mechanisms would provide the same results for the doubly OZI disconnected processes. However, difficulty will arise due to possible contributions from a large number of intermediate processes. A further reasonable assumption then is to consider the dominant

processes based on the available experimental information. As shown by PDG [22], J/ψ has the largest partial decay widths to $\rho\pi$, and ω meson has also strong couplings to $\rho\pi$. Meanwhile, for these f_0 states, the decay of $f_0^i \rightarrow \pi\pi$ is one of the most important channels, e.g. $f_0(1370) \rightarrow \pi\pi$. This makes the $\rho\pi$ intermediate exchange a leading contribution to $J/\psi \rightarrow \omega f_0^3$. Similarly, the intermediate $K^*\bar{K} + c.c.$ exchange should be also a dominant contribution to $J/\psi \rightarrow \phi f_0^1$: the partial decay width of $J/\psi \rightarrow K^*\bar{K} + c.c.$ is about one order of magnitude larger than $J/\psi \rightarrow \phi f_0^1$; the ϕ meson couples strongly to K^*K ; and $K\bar{K}$ is the dominant decay channel for the $f_0(1710)$. Furthermore, note that the other hadronic decay channels of J/ψ are almost the same order of magnitude as $J/\psi \rightarrow \omega f_0(1710)$, or just one magnitude larger than $J/\psi \rightarrow \phi f_0(1710)$. Their rescatterings to final state $V f_0^i$ would be strongly suppressed due to their weak couplings to $V f_0^i$ and energy suppressions at the rescattering vertices.

Following the meson exchange mechanisms, we calculate the meson loops in Fig. 4 as a leading contribution to the doubly disconnected process. Supposing the OZI violation effects are from the intermediate meson exchange processes, we can estimate the meson exchange contribution and compare it with the “tree” diagram for $J/\psi \rightarrow V\bar{P} + c.c.$, e.g. $J/\psi \rightarrow K^*\bar{K} + c.c.$ and $J/\psi \rightarrow \rho\pi + c.c.$ Since the $J/\psi V\bar{P}$ vertices are the same between the tree and loop transitions, the ratio between the meson-exchange loop and the tree process of $J/\psi \rightarrow V\bar{P} + c.c.$ will then highlight the OZI violations in this dynamical process. For the purpose of gaining information about the rough behavior of the energy-evolution of the doubly OZI processes, the energy-evolution of this ratio is sufficient to account for the energy-dependence of the doubly OZI violation transitions.

The transition amplitude can be written as

$$M_{fi} = -i \int \frac{d^4 p_2}{(2\pi)^4} T_v^\beta \left(g_\beta^\lambda - \frac{p_{1\beta} p_1^\lambda}{p_1^2} \right) T_{0\lambda} T_s \frac{F(p_2)}{a_1 a_2 a_3} \delta^4(p_0 - p_v - P_s), \quad (10)$$

where the vertex functions are

$$\begin{aligned} T_v^\beta &= i \frac{g_v}{M_v} \epsilon^{\mu\nu\alpha\beta} p_{v\mu} \epsilon_{f\nu} p_{2\alpha}, \\ T_\lambda &= i \frac{g_0}{M_0} \epsilon_{\lambda\sigma\tau\delta} p_0^\sigma \epsilon_i^\tau p_3^\delta, \\ T_0 &= i g_s M_s, \end{aligned} \quad (11)$$

and $a_1 = p_1^2 - m_1^2 + i\epsilon$, $a_2 = p_2^2 - m_2^2 + i\epsilon$, and $a_3 = p_3^2 - m_3^2 + i\epsilon$ are the denominators of the meson propagators. The coupling constants g_v , g_s and g_0 can be determined independently in meson decays. For the J/ψ hadronic decays, large branching ratios for $J/\psi \rightarrow K^*\bar{K} + c.c.$ and $\rho\pi + c.c.$ imply that the meson loop transitions may have significant contributions to those decays where the final state mesons have also large couplings to the exchanged mesons. This is indeed the case for $J/\psi \rightarrow V f_0(1710)$, where the couplings for ϕK^*K , ωK^*K , and $f_0(1710)K\bar{K}$ are significantly large. Therefore, this simple argument will allow us to consider only the dominant meson exchange loops in the calculations.

In the above equation, the coupling g_0 can be determined by the decays of $J/\psi \rightarrow K^*\bar{K} + c.c.$ [Fig. 4(a) and (b)], or $J/\psi \rightarrow \rho\pi + c.c.$ [Fig. 4(c)], of which large branching ratios are observed in experiment, e.g.

$$g_0^2 = \frac{12\pi M_0^2}{|\mathbf{p}_1|^3} \Gamma_{J/\psi \rightarrow V\bar{P}}^{exp}, \quad (12)$$

where $\Gamma_{J/\psi \rightarrow K^*\bar{K} + c.c.}^{exp} = (9.2 \pm 0.8) \times 10^{-3}$ and $\Gamma_{J/\psi \rightarrow \rho\pi + c.c.}^{exp} = (1.27 \pm 0.09)\%$ are from the estimate of Particle Data Group [22].

For $f_0 \rightarrow K\bar{K}$ in the f_0 c.m. system, the coupling constant can be derived via

$$g_s^2 = \frac{8\pi}{|\mathbf{k}|} \Gamma_{f_0 \rightarrow K\bar{K}}^{exp}, \quad (13)$$

which is consistent with the studies of $f_0 \rightarrow PP$ in the determination of the mixing matrix elements [15]; $|\mathbf{k}|$ is the magnitude of the three momenta carried by the final-state kaon (anti-kaon). With the estimate of $b.r. f_0 \rightarrow K\bar{K} = 0.60$ [15], and $\Gamma_T(1710) = 140$ MeV [22], we have $\Gamma_{f_0 \rightarrow K\bar{K}}^{exp} = 84$ MeV; the coupling g_s can then be determined.

We determine the VVP couplings, i.e., $g_{\phi K^*K}$, $g_{\omega K^*K}$, and $g_{\omega\rho\pi}$, in the SU(3)-flavor-symmetry limit. Since ω and ϕ are almost ideally mixed, the SU(3) symmetry leads to

$$g_{\omega\rho\pi} = 3g_{\omega\rho^0\pi^0}, \quad g_{\phi K^*K} = 2\sqrt{2}g_{\omega\rho^0\pi^0}, \quad g_{\omega K^*K} = 2g_{\omega\rho^0\pi^0}, \quad (14)$$

where $g_{\omega\rho^0\pi^0}^2 \simeq 84$ is determined in vector meson dominance (VMD) model in $\omega \rightarrow \pi^0 e^+ e^-$ [22].

We apply the Cutkosky rule to the loop calculation, where the mesons from the decay vertex are assumed to be on-shell, i.e. $p_1^2 = m_1^2$ and $p_3^2 = m_3^2$. The transition amplitude then reduces to

$$M_{fi} = -\frac{ig_0g_vg_s}{64\pi^2}[p_v \cdot p_0\epsilon_f \cdot \epsilon_i - p_v \cdot \epsilon_i p_0 \cdot \epsilon_f] \frac{M_s|\mathbf{p}_3|}{M_0^2M_v} \mathcal{S}, \quad (15)$$

where

$$\mathcal{S} = \int d\Omega \frac{(p_s - p_3)^2 F((p_s - p_3)^2)}{(p_s - p_3)^2 - m_2^2}, \quad (16)$$

is the reduced loop integral after applying the on-shell condition for the integrand; The momentum conservation, $p_2 = p_s - p_3$, hence leads to the angular dependence of the integrand in the above equation since $p_2^2 = (p_s - p_3)^2 = M_s^2 + m_3^2 - 2E_sE_3 + 2|\mathbf{p}_s||\mathbf{p}_3|\cos\theta$. The form factor $F((p_s - p_3)^2)$ is included to take into account the off-shell effects for the exchanged meson in the final state interactions.

In the loop transition, the polarizations of the initial-final state vector mesons, longitudinal-transverse and vice versa (TL), will not contribute to the integrals. Also, polarization of longitudinal-longitudinal (LL) vanishes in the tree transitions. We hence take the transverse-transverse (TT) ratio between the loop and tree transitions to define

$$r_c \equiv -\frac{g_vg_s}{64\pi^2} \frac{M_sE_v}{M_0M_v} \mathcal{S}, \quad (17)$$

where E_v is the energy of the final state vector meson, and r_c reflects the energy evolution of the meson loop transitions. For the loop transition, the LL component compared to the LL ones is M_v/E_v suppressed at higher energies, i.e., $M_0 \rightarrow \infty$. Therefore, r_c is a good quantity for describing the energy evolutions of the loop transitions relative to the tree ones. A feature arising from r_c is that $E_v/M_0 \rightarrow 1/2$ in the limit of $M_0 \rightarrow \infty$. Therefore, for sufficiently large M_0 , the energy evolution of r_c will be determined by $\mathcal{S}$.

The meson loop integration of Eq. 16 can be rewritten as

$$\mathcal{S} = -2\pi \frac{B}{A} \int d\cos\theta \frac{1 + A\cos\theta}{1 + B\cos\theta} F((p_s - p_3)^2), \quad (18)$$

where

$$\begin{aligned} A &\equiv 2|\mathbf{p}_s||\mathbf{p}_3|/(M_s^2 + m_3^2 - 2E_sE_3) \\ B &\equiv 2|\mathbf{p}_s||\mathbf{p}_3|/(M_s^2 + m_3^2 - 2E_sE_3 - m_2^2). \end{aligned} \quad (19)$$

For the following situations with explicit considerations of the form factors, the integration can be worked out analytically:

i) Without form factor: $F((p_s - p_3)^2) = 1$

The integration gives

$$\mathcal{S}_a = 2\pi \left[2 - \frac{B-A}{AB} \ln \frac{1-B}{1+B} \right]. \quad (20)$$

ii) Monopole form factor: $F((p_s - p_3)^2) = (\Lambda_1^2 - m_2^2)/(\Lambda_1^2 - p_2^2)$

Define

$$\begin{aligned} C &\equiv 2|\mathbf{p}_s||\mathbf{p}_3|/(\Lambda_1^2 - (M_s^2 + m_3^2 - 2E_sE_3)) \\ \lambda &\equiv (\Lambda_1^2 - m_2^2)/(\Lambda_1^2 - (M_s^2 + m_3^2 - 2E_sE_3)), \end{aligned} \quad (21)$$

we have

$$\mathcal{S}_b = 2\pi\lambda \left[\frac{1}{C} \ln \frac{1+C}{1-C} + \frac{A-B}{A(B-C)} \ln \frac{(1-B)(1+C)}{(1+B)(1-C)} \right]. \quad (22)$$

iii) Dipole form factor: $F((p_s - p_3)^2) = [(\Lambda_2^2 - m_2^2)/(\Lambda_2^2 - p_2^2)]^2$

The integration gives

$$\mathcal{S}_c = 2\pi\lambda^2 \left\{ \frac{2}{1-C^2} + \frac{A-B}{A} \left[\frac{2C}{(B-C)(1-C^2)} + \frac{B}{(B-C)^2} \ln \frac{(1-B)(1+C)}{(1+B)(1-C)} \right] \right\}. \quad (23)$$

In (ii) and (iii) parameters Λ_1 and Λ_2 have a range of $1 \sim 2$ GeV, which are commonly adopted in the literature [23]. In general, the adopted value for Λ_1 is smaller than that for Λ_2 . In this calculation, we adopt $\Lambda_1 = 1.2$ GeV, and $\Lambda_2 = 1.9$ GeV, since they lead to the same results for $\mathcal{S}_b$ and $\mathcal{S}_c$ in J/ψ decays. If the same value for Λ_1 and Λ_2 is adopted, we find $\mathcal{S}_c/\mathcal{S}_b \simeq 0.33 \sim 0.55$ within the range of $1.2 \sim 1.9$ GeV for the cut-off energy.

In Table I, the ratio r_c is calculated for $J/\psi \rightarrow K^*\bar{K} + c.c. \rightarrow \phi f_0^i$. Without form factor, the loop transition accounts for large fraction of the $J/\psi \rightarrow K^*\bar{K} + c.c.$ decay, and obviously overestimates the branching ratios for $J/\psi \rightarrow V f_0^i$. In contrast, the inclusion of form factors will significantly suppress the loop transition strengths. Roughly to say, the loop transition has the largest contributions to the production of $f_0(1710)$. This can be explained by the large $f_0(1710)K\bar{K}$ couplings. Interestingly, the K^*K loop contributions to the branching ratio of $J/\psi \rightarrow \phi f_0(1710) \rightarrow \phi K\bar{K}$ have the same order of magnitude as the experimental data. This directly shows that large OZI violations occur via the loop transitions. Also, the suppressed loop contributions in the production of $f_0(1500)$ and $f_0(1370)$ are consistent with the experimental results [1] and calculations presented in Ref. [15].

For $J/\psi \rightarrow \omega f_0^i$, in addition to the K^*K loop transition, the $\rho\pi$ loop may also be significantly large. In Table II, the loop contributions from $J/\psi \rightarrow K^*\bar{K} + c.c. \rightarrow \omega f_0^i$ are presented. Again, we obtain a relatively larger branching ratio for the $f_0(1710)$ than for the $f_0(1500)$ and $f_0(1370)$. Compared with the experimental data [2], where $\Gamma_{J/\psi \rightarrow \omega f_0(1710)} > \Gamma_{J/\psi \rightarrow \phi f_0(1710)}$, it shows that the loop transition contributes slightly larger in $J/\psi \rightarrow \phi f_0(1710)$ than in $J/\psi \rightarrow \omega f_0(1710)$. This originates from $g_{\phi K^*K} > g_{\omega K^*K}$ in the loop.

The story changes in the $\rho\pi$ loop transition. As presented in Table III, this process has sizeable contributions to the $f_0(1500)$ and $f_0(1370)$ productions, but is negligibly small in the production of $f_0(1710)$. This suggests much more important roles are played by the $\rho\pi$ loop transitions for those two states.

Although we do not present the estimate of the contributions from the $\rho\pi$ loop transition to $J/\psi \rightarrow \rho\pi + c.c. \rightarrow \phi f_0^i$, it is worth noting that the $\phi\rho\pi$ coupling should not be small. As quoted by PDG [22], the branching ratio for $\phi \rightarrow \rho\pi + \pi^+\pi^-\pi^0$ is about 15.4%. Supposing a large fraction of the decays is via $\phi \rightarrow \rho\pi$, it will give a large coupling, and naturally lead to large branching ratios for the $f_0(1370)$ and $f_0(1500)$ in the $\phi\pi\pi$ channel.

It should be noted that the loop transitions could contribute more to the branching ratios. Due to the on-shell approximation applied, we have only included the imaginary part of the transition amplitudes in the calculation, and neglected contributions from the real part. More rigorous approaches would also consider the latter via e.g. disperse relation [24]. For the purpose of illustrating large OZI violations via the loop transitions, and investigating their energy evolution, it is sufficient to only consider the imaginary amplitudes from the on-shell approximation.

To relate the above loop transitions to the OZI violation parameter r , we first note that the ratio r does not equivalent to r_c . In the definition of r_c , the tree coupling is given by $J/\psi \rightarrow K^*\bar{K} + c.c.$, while for r , the tree coupling is from the singly disconnected processes. Due to lack of constraints on the singly disconnected processes of $J/\psi \rightarrow \phi f_0$, e.g. Fig. 1(a) and (b), we cannot directly derive r . However, notice that the major difference between r and r_c is the different coupling at the J/ψ decay vertex. We can then estimate:

$$\frac{r(M_\Upsilon)}{r(M_{J/\psi})} \simeq \frac{r_c(M_\Upsilon)}{r_c(M_{J/\psi})}. \quad (24)$$

Such a relation must be a robust one since we do not include all the intermediate meson exchange contributions. However, as we addressed in the previous part, the dominant meson-loop contributions are from the processes that the couplings at those three vertices are strong. In this sense, the neglect of other meson exchange contributions should not result in magnitude uncertainties in the above estimate. As confirmed by the calculations that the loop transitions are the dominant contributions to the OZI violation processes and have the same order of magnitude as the singly disconnected processes, we adopt $r(M_{J/\psi}) = 2.2$ from Ref. [15] to derive $r(M_\Upsilon) \simeq 0.26 \sim 0.28$, which indicates that the loop transition amplitude is indeed suppressed at the Υ mass region. The variation is from calculations for the K^*K and $\rho\pi$ loops, respectively.

Mapping $r(M_\Upsilon) \simeq 0.27 \pm 0.01$ to Fig. 2 (see the arrows in the figure), we find that $R_1^{OZI} \simeq 1.67 \pm 0.04$ in contrast with $R_1^{OZI} = 0.15$ at $r(M_{J/\psi}) = 2.2$. As shown by the curves in Fig. 2 between $r = 0.27 - 2.2$, R_i^{OZI} experiences a dramatical change, which suggests a measurable effect from the possible OZI violation processes in the energy evolutions. In the small r region, where the OZI rule applies, the production widths for $f_0(1500)$ and $f_0(1370)$ in the ϕ channel will be significantly suppressed in comparison with the ω channel. At the Υ energy, i.e. $r \simeq 0.27$, the ratios R_2^{OZI} and R_3^{OZI} are also inversely changed in contrast with the values derived at the J/ψ mass region.

The energy evolution of r_c in those three loops are presented in Fig. 5. Exponential drop can be seen at small M_0 region, and it slightly levels off above 7 GeV. This indicates explicitly that the contributions from the loop transitions will decrease much faster than the tree transitions. This is consistent with the qualitative analyses made earlier. One can also see that the three r_c 's have similar behaviors with the increasing energies. This eventually justifies the above simple relation as a general results for relating r to r_c through the dominant loop transitions.

Undoubtedly, the measurement of R_i^{OZI} in the Υ decays also examines the glueball- $Q\bar{Q}$ mixing scheme proposed for the scalar mesons. As presented in Fig. 2, the dramatic change of the relative positions of R_i^{OZI} for these three f_0 states

also depend on the mixing matrix elements. For example, in addition to the measurement of R_1^{OZI} for the $f_0(1710)$ in $\Upsilon \rightarrow V f_0(1710) \rightarrow V K \bar{K}$, it is also interesting to measure R_3^{OZI} for the $f_0(1370)$ in $\Upsilon \rightarrow V f_0(1370) \rightarrow V \pi \pi$. The values for R_1^{OZI} and R_3^{OZI} if match the pattern of Fig. 2, would be a strong support for the glueball- $Q\bar{Q}$ mixings. The R_2^{OZI} for the $f_0(1500)$ can also be investigated though difficulty may arise from the small branching ratios for both $\Upsilon \rightarrow \phi f_0(1500)$ and $\Upsilon \rightarrow \omega f_0(1500)$. Although, high statistics in experiment should be required to isolate these two decay channels, the energy-evolution trends of R_i^{OZI} seem to provide a way to disentangle the correlation between the OZI-rule violation and the structure of those scalar mesons.

In summary, we discussed a possible way to determine the scalar structures by clarifying the role played by the OZI-rule violation. The latter was correlated with the glueball- $Q\bar{Q}$ mixings in $J/\psi \rightarrow V f_0^i$. Since the flavor wavefunctions for ω and ϕ are almost ideally mixed, the decay channels into ω and ϕ in association with the scalar mesons, respectively, serve as a flavor filter for probing the $Q\bar{Q}$ contents of the scalars. This allows us to separate out the doubly OZI disconnected processes, of which the effects can be measured by the branching ratio fractions between ϕf_0^i and ωf_0^i , i.e. R_i^{OZI} . Since the energy evolution of R_i^{OZI} is mostly determined by the energy evolution of the doubly disconnected processes relative to the singly disconnected ones, the suppression of the doubly disconnected process at higher energies, e.g. in Υ decays, will lead to dramatic changes to R_i^{OZI} with certain patterns. Observation of such a change will provide direct information about the scalar meson structures.

Although at this moment, experimental data for $\Upsilon \rightarrow V f_0^i$ are unavailable, clear signals for $f_0(1710)$ have been reported in $\Upsilon \rightarrow \gamma f_0(1710) \rightarrow \gamma K \bar{K}$ [22]. With an increasing statistics, one may have access to the above mentioned channels at future facilities for heavy quarkonium physics [25, 26].

The authors thank F.E. Close for many useful discussions and comments on this work. This work is supported, in part, by grants from the U.K. Engineering and Physical Sciences Research Council Advanced Fellowship (Grant No. GR/S99433/01), and the Institute of High Energy, Chinese Academy of Sciences.

-
- [1] M. Ablikim *et al.* [BES Collaboration], arXiv:hep-ex/0411001.
 - [2] M. Ablikim *et al.* [BES Collaboration], Phys. Lett. B **603**, 138 (2004) [arXiv:hep-ex/0409007].
 - [3] M. Ablikim *et al.* [BES Collaboration], Phys. Lett. B **598**, 149 (2004) [arXiv:hep-ex/0406038].
 - [4] S. Jin, Plenary talk at 32nd International Conference on High-Energy Physics (ICHEP2004), Beijing (2004).
 - [5] F.E. Close and N.A. Tornqvist, J. Phys. G **28**, R249 (2002).
 - [6] F.E. Close and C. Amsler, Phys. Lett. B **353**, 385 (1995); Phys. Rev. **D53**, 295 (1996).
 - [7] F.E. Close, Plenary talk at 32nd International Conference on High-Energy Physics (ICHEP2004), Beijing (2004) [archive:hep-ph/0411396].
 - [8] C. Morningstar and M. Peardon, Phys. Rev. D **56**, 4043 (1997).
 - [9] G. Bali *et al.*, UKQCD Collaboration, Phys. Lett. B **309**, 378 (1993).
 - [10] M.S. Chanowitz, Proc. 6th Int. Workshop on $\gamma\gamma$ Collisions, Edited by R. Lander (World Scientific, Singapore, 1984).
 - [11] S. Narison, Nucl. Phys. B **509**, 312 (1998).
 - [12] F.E. Close, G.R. Farrar, and Z. Li, Phys. Rev. D **55**, 5749 (1997).
 - [13] D.V. Bugg, M. Peardon, and B.S. Zou, Phys. Lett. B **486**, 49 (2000).
 - [14] F.E. Close and A. Kirk, Phys. Lett. B **483**, 345 (2000).
 - [15] F.E. Close and Q. Zhao, Phys. Rev. D **71**, 094022 (2005) [arXiv:hep-ph/0504043]; Phys. Lett. B **586**, 332 (2004) [arXiv:hep-ph/0402090].
 - [16] S. Okubo, Phys. Lett. **5**, 1975 (1963); G. Zweig, in *Development in the Quark Theory of Hadrons*, edited by D.B. Lichtenberg and S.P. Rosen (Hadronic Press, Massachusetts, 1980); J. Iizuka, Prog. Theor. Phys. Suppl. **37**, 38 (1966).
 - [17] A. Seiden, H. F.-W. Sadrozinski, and H. E. Haber, Phys. Rev. D **38**, 824 (1988).
 - [18] H.J. Lipkin, Phys. Rev. Lett. **13**, 590 (1964); **14**, 513 (1965); Phys. Rep. **8C**, 173 (1973); Nucl. Phys. **B244**, 147(1984); **B291**, 720 (1987); Phys. Lett. **B179**, 278 (1986).
 - [19] P. Geiger and N. Isgur, Phys. Rev. D **47**, 5050 (1993).
 - [20] H.J. Lipkin and B.S. Zou, Phys. Rev. D **53**, 6693 (1996).
 - [21] M. P. Locher, Y. Lu and B. S. Zou, Z. Phys. A **347**, 281 (1994) [arXiv:nucl-th/9311021].
 - [22] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
 - [23] X.Q. Li, D.V. Bugg, and B.S. Zou, Phys. Rev. D **55**, 1421 (1997).
 - [24] H.-Y. Cheng, C.-K. Chua, and A. Soni, Phys. Rev. D **71**, 014030 (2005).
 - [25] S.A. Dytman *et al.* [CLEO Collaboration], hep-ex/0307035.
 - [26] N. Brambilla *et al.*, hep-ph/0412158.
 - [27] Here, we neglect the trivial kinematic dependence arising from the mass differences between ϕ and ω .

	$f_0(1710)$			$f_0(1500)$			$f_0(1370)$		
	(i)	(ii)	(iii)	(i)	(ii)	(iii)	(i)	(ii)	(iii)
$r_c(M_{J/\psi})$	0.550	0.163	0.165	0.196	0.056	0.057	0.149	0.041	0.042
$b.r.(J/\psi \rightarrow \phi f_0)$	19.31	1.71	1.73	2.95	0.24	0.24	1.83	0.14	0.15
$b.r.(J/\psi \rightarrow \phi f_0 \rightarrow \phi K \bar{K})$	11.59	1.02	1.04	2.53	0.02	0.02	0.04	0.00	0.00
$b.r.(exp)$	(2.0 ± 0.7)			(0.8 ± 0.5)			(0.3 ± 0.3)		

TABLE I: The intermediate K^*K meson exchange contributions to the decay of $J/\psi \rightarrow \phi f_0 \rightarrow \phi K \bar{K}$ for three cases: (i) *without* form factors; (ii) *with* monopole form factors; and (iii) *with* dipole form factors. The channel to which the loop transition has significant contributions is highlighted by bold characters. The numbers in the round brackets are data for $J/\psi \rightarrow \phi f_0 \rightarrow \phi K \bar{K}$ from BES [1]. The branching ratios are in a unit of 10^{-4} .

	$f_0(1710)$			$f_0(1500)$			$f_0(1370)$		
	(i)	(ii)	(iii)	(i)	(ii)	(iii)	(i)	(ii)	(iii)
$r_c(M_{J/\psi})$	0.479	0.139	0.141	0.172	0.047	0.049	0.131	0.034	0.036
$b.r.(J/\psi \rightarrow \omega f_0)$	16.47	1.39	1.43	2.39	0.18	0.19	1.46	0.10	0.11
$b.r.(J/\psi \rightarrow \omega f_0 \rightarrow \omega K \bar{K})$	9.88	0.84	0.86	0.21	0.02	0.02	0.03	0.00	0.00
$b.r.(exp)$	(13.2 ± 2.6)			...			...		

TABLE II: The intermediate K^*K meson exchange contributions to the decay of $J/\psi \rightarrow \omega f_0 \rightarrow \omega K \bar{K}$ presented in a way similar to Table I. The numbers in the round brackets are data for $J/\psi \rightarrow \omega f_0 \rightarrow \omega K \bar{K}$ from BES [2], and the dots denote the unavailability of the experimental data. The branching ratios are in a unit of 10^{-4} .




FIG. 1: Schematic processes for the production of f_0^i in $J/\psi \rightarrow \phi f_0^i$.

	$f_0(1710)$			$f_0(1500)$			$f_0(1370)$		
	(i)	(ii)	(iii)	(i)	(ii)	(iii)	(i)	(ii)	(iii)
$r_c(M_{J/\psi})$	0.236	0.072	0.078	0.494	0.143	0.156	0.565	0.159	0.175
$b.r.(J/\psi \rightarrow \omega f_0)$	5.26	0.49	0.57	25.87	2.18	2.59	35.84	2.85	3.43
$b.r.(J/\psi \rightarrow \omega f_0 \rightarrow \omega \pi \pi)$	0.35	0.03	0.04	9.03	0.76	0.90	7.17	0.57	0.69
$b.r.(exp)$		...			...			...	

TABLE III: The intermediate $\rho\pi$ meson exchange contributions to the decay of $J/\psi \rightarrow \omega f_0 \rightarrow \omega \pi \pi$ presented in a way similar to Table I. The dots denote the unavailability of the experimental data. The branching ratios are in a unit of 10^{-4} .




FIG. 2: Energy evolution of R_i^{OZI} in terms of parameter r .




FIG. 3: The OZI doubly disconnected processes in terms of intermediate virtual meson exchanges.




FIG. 4: The OZI violations from intermediate K^*K and $\rho\pi$ meson exchanges as dominant contributions to the doubly disconnected processes.




FIG. 5: The energy evolution of the ratio between the dominant meson loop transitions and the corresponding tree transitions. The solid curve is for K^*K loop in the $\phi f_0(1710)$ channel, while the dashed and dotted curves are for the K^*K and $\rho\pi$ loops in the $\omega f_0(1710)$ channels.